

# Evaluation of Using Bioactivator

*by Hajas Article*

---



**Submission date:** 21-Apr-2022 04:21PM (UTC+0700)

**Submission ID:** 1816220352

**File name:** 18527-Layout\_revisi\_2.docx (106.43K)

**Word count:** 2533

**Character count:** 14060



# The Evaluation of Using Bio Activators Based on Rumen Content of Dry Matter, Organic Matter, and Crude Protein on Oil Palm Fronds as Ruminant Feed

Tri Astuti<sup>a</sup>, Syahro Ali Akbar<sup>a</sup>, Delsi Afrini<sup>b</sup>, M. Nasir Rofiq<sup>c</sup>, Fajri Basyirun<sup>d</sup>

<sup>a</sup>Department of Animal Science, Faculty of Agriculture, University of Mahaputra Muhammad Yamin, West Sumatra, Indonesia

<sup>b</sup>Department of Agribusiness, Faculty of Agriculture, University of Mahaputra Muhammad Yamin, West Sumatra, Indonesia

<sup>c</sup>BPPT Serpong, Indonesia

<sup>d</sup>Department of Economic Education, Faculty of Education, University of Mahaputra Muhammad Yamin, West Sumatra, Indonesia

\*Corresponding author: E-mail: [adektuti@gmail.com](mailto:adektuti@gmail.com)

## ARTICLE INFO

## ABSTRACT

### Article history:

Submission: 29-10-2021

Accepted: 07-04-2022

Published:

This study aims to examine the effect of using rumen content bioactivators with the addition of different energy sources on nutrient content, fiber fraction content, and energy estimation in palm leaves and midribs that will be used as feed ingredients for ruminants. Factorial completely randomized design 2x4 with 4 replications for each treatment was used in this study. Factor A is the source of the rumen contents, where A1 = Cattle rumen, A2 = Buffalo rumen. Factor B is a microbial energy supply; B1 = Molasses, B2 = Molasses + palm frond extract, B3 = Molasses + palm leaf extract, B4 = Molasses + palm frond and leaf extract. Parameters observed were dry matter content, organic matter, crude protein. The results of the variance analysis showed that there was no interaction (P > 0.05) between the source of rumen content and the type of microbial energy supply. The dry matter content of palm fronds is not affected by the source of the rumen content of cows or buffaloes. The source of the rumen content based on cattle or buffaloes has no effect on the dry matter content of palm fronds.

Keywords: Bioactivator, oil palm frond, ruminant.

## INTRODUCTION

Feeding is the main component that for the success of a livestock business, and the feeding is the highest cost of all the daily operational of a livestock business [1, 2, 3]. The Grass as the main source of forage for ruminants [4, 5, 6, 7] is currently limited in availability due to the conversion of agricultural land into housing and other infrastructure development. This causes an effort to find alternative sources as a substitute for forage in ruminants. Palm oil plantations in addition to providing benefits from their production [8] and processing for food

[9], there is also a large amount of waste [10], both in terms of quantity and type, both from plantation waste and waste from processing factories. Oil palm fronds is solid waste from oil palm plantations that have the potential to be used as basal feed for ruminants [11, 12, 13]. The oil palm fronds must be trimmed to reach 8.6 tons per hectare/year [11]. Utilization of oil palm fronds is a potential as substitution of field grass for the source of forage ruminant feed [14], because of its abundant availability and is not seasonal. Palm oil by-products can be pursued as potential feed for ruminants. The utilization of plantation waste as feeding is always constrained by the presence of lignocellulosic bonds which usually have high lignin content and low nutritional value.[15] that oil palm fronds contain 3.25% crude protein, 49.51% dry matter, and 95.96% organic matter. [16] the lignin content of oil palm fronds is 15.34%. In addition, there are many thorns and sticks of palm leaves on the oil palm fronds which are quite sharp, difficult to digest, and it is feared that it will be risky for livestock if consumed in the long term. It needs to treat feed processing technology before using a feeding. Treatment of feed ingredients through a biotechnology fermentation approach is considered easier, cheaper, and environmentally friendly compared to chemical treatment [17]. Astuti et. al [18] showed that the total number of colonies of local microorganism (MOL) bacteria in the rumen with the addition of energy sources (tofu soaking water and molasses) and palm fronds was  $10 \times 10^{10}/\text{ml}$ . There are so many waste rumen of cattle and buffalo from the slaughterhouse. In 2018 the cattle and buffalo slaughtered at slaughterhouses was 1,147,657 [19]. The Slaughter of ruminants is always increasing every time, in line with the increasing human needs for ruminant meat food. Rumen fluid also contains the enzymes  $\alpha$ -amylase, galactosidase, hemicellulase, cellulase, and xylanase [20]. Oil palm fronds can be used as a source of forage feed ingredients by improving nutrient quality both through physical and biological processing using rumen content bio activators. Fermentation process oil palm frond with bio activator cattle rumen causes a decrease in crude protein by 3.93% and increase crude fiber by 42.96% [21]. The composition of amino acids, minerals, and vitamins as well as enzymes, also depends on the treatment of the feed given [22]. Rumen fluid enzymes as an alternative technology that can be used to hydrolyze crude fiber and increase the nutritional value of local feed raw materials [23]. This study aimed to evaluate the nutrient content, fiber fraction, and energy estimation of palm fronds incubated with bio activators from the rumen contents of cattle and buffalo. Microorganisms contained in the contents of the rumen are "bred" first by feeding/supplying the required energy that is adjusted to the expected target of enzyme production.

## MATERIALS AND METHODS

This study was conducted to determine the effectiveness of the use of rumen content bio-activator on oil palm fronds.

### Bio-Activator Production Process

The rumen contents were taken from the slaughterhouse. The contents of the rumen are taken using a thermos and filled with hot water to keep the temperature stable which consisted of beef rumen contents and buffalo rumen contents, each mixed with soybean soaking water, molasses in a certain ratio as a source of protein and carbon which is the rumen

microbial energy supply. In addition, a substrate is also added as a source of enzymes that will be expected to be produced by rumen microbes. There are 4 (four) types of bio activators with the addition of different enzyme sources. This solution mixture was incubated for 7 days.

### Oil Palm Fronds Fermentation Process

The fronds and palm leaves are mashed then fermented with rumen contents bio-activator for 7 days. The factorial randomized design used was a 2 x 4 with 4 replications for each treatment. Factor A was the type of rument content: A1 = Cow rument, A2 = Buffalo rument. Factor microbial energy supply; B1 = Molasses, B2 = Molasses + palm frond extract, B3 = Molasses + palm leaf extract, B4 = Molasses + palm frond and leaf extract. Parameters measured were dry matter content, organic matter, crude protein.

### Statistical Analysis

All data tested by analysis variant and the significantly affect data were further tested by Duncan's multiple range test.

### RESULTS AND DISCUSSIONS

The average dry matter, organic matter, and crude protein content of oil palm fronds incubated with rumen bio-activator.

The data in table 1 was the analysis of the average dry matter content, organic matter, and crude protein of palm fronds incubated with cow rumen content bio-activator and buffalo rumen content with the addition of different enzyme energy sources.

The results of the diversity analysis showed that there was no interaction ( $P > 0.05$ ) between the source of rumen content and the type of microbial energy supply. Sources of rumen content of cows and buffaloes gave no different effect on the dry matter content of oil palm fronds, as well as the addition of energy and feed sources for microbes gave no different effects as well. This is because in this study the contents of the rumen used were rumen fluid which contained a lot of rumen microbes. Some microorganisms digest glucose and produce water, carbon dioxide, and large amounts of energy (ATP) which are used for growth. The results of the analysis of diversity showed that the organic matter content in this study showed no significant effect ( $P > 0.05$ ), on the content of oil palm fronds incubated using cow rumen content bio-activator and buffalo rumen content (92.57 vs 92.56 %). This shows that the activities carried out by cow rumen microbes and buffalo rumen do not differ in degrading organic matter found in oil palm midribs. There was a significant interaction ( $P < 0.05$ ) between the source of rumen content and the source of microbial energy on the crude protein content of the palm fronds. This is because in the contents of the rumen there are various rumen microbes that grow and develop based on the energy source provided. Microbes (bacteria, and fungi) are proteins. The best crude protein content was found in palm fronds incubated with cattle rumen contents with the addition of molasses, soybean soaking water, and palm leaves extract (9.03%). Based on Table 1, it can be seen that the average protein content was higher in palm fronds fermented using cow rumen content bio activator compared to buffalo rumen content (8.14 vs 8.01%).

Table 1. The Average Nutrient Content of Oil Palm Frond Incubation Using Bio-Activator of Cattle and Buffalo Rumen Content

Factor A	Factor B				Average
	B1	B2	B3	B4	
Dry Matter Content					
A1	36.49	35.33	36.14	36.37	36.08
A2	40.43	42.63	42.20	44.66	42.48
Average	38.46	38.98	39.17	40.52	
Organic Matter Content					
A1	92.54	92.81	92.42	92.50	92.57
A2	92.64	92.81	92.39	92.40	92.56
Average	92.59	92.81	92.40	92.47	
Crude Protein Content					
A1	8.04 <sup>aA</sup>	7.35 <sup>bA</sup>	9.03 <sup>aA</sup>	8.15 <sup>aA</sup>	8.14
A2	7.81 <sup>bA</sup>	8.09 <sup>aA</sup>	7.19 <sup>bA</sup>	8.98 <sup>aA</sup>	8.01
Average	7.92	7.72	8.11	8.98	

Note: The superscripts (a,b) in the same row and (A,B) in the same column showed significantly different effects (P<0.05)

Based on the results of the study, it was also seen that the average crude protein content based on the source of enzyme supply was the highest in the treatment of rumen contents mixed with molasses, soybean soaking water, leaves, and palm fronds (8.98%). This crude protein content is higher than the research by [15] which found that the crude protein content of fermented oil palm fronds with was 5.35%. When compared with the protein content of palm midrib without treatment, there was an increase of 121.23 – 177.85% in this study (3.25 vs 7.19-9.03%). This was because the rumen contents bio activator consists of bacteria and fungi whose main component is protein. Riswandi [24] stated that as the number of microbes increases, the crude protein content of fermented feed will increase, because microbes are a source of single-cell protein. In addition, in the process of making bio-activators, the rumen contents are mixed with soy water immersion, where the crude protein content of soybeans is 46.2%.

10

## CONCLUSIONS

Based on data, it can be concluded that there was no interaction effect on dry matter and organic matter content of oil palm fronds by bio-activator treatment, and there is a highly significant interaction effect on crude protein content.

## REFERENCES

- [1] S. Said, "Integrated Livestock Business and Industry in Indonesia", IOP Conf. Ser.: Earth Environ. Sci., 465, 012003, pp. 1-6, 2020. doi:10.1088/1755-1315/465/1/012003

- [2] L. Lindawati, D. Henderawan, I. Zulfida, and K. N. Lumbantoruan, "Improvement Strategy of Livestock Business in Deli Serdang", IOP Conf. Series: Earth Environ. Sci., 454, 012024, pp. 1-6, 2020. doi:10.1088/1755-1315/454/1/012024
- [3] N. M. Santa, F. N. Sompie, and W. Waworundeng, "Livestock Business Development of Border Areas in North Sulawesi Province", IOP Conf. Series: Earth Environ. Sci., 473, 012139, pp. 1-4, 2020. doi:10.1088/1755-1315/473/1/012139
- [4] K. Muatip, T. Widiyastuti, N. N. Hidayat, H. Purwaningsih, E. Purwanto, and G. G. Setya, "Forage Business at Kebumen District Central Java Province", Animal Production, Vol. 19, no. 2, pp. 135-142, 2017.
- [5] R. Malaka, R. Islamiyati, S. Baco, A. Sabir, Hamsah, and G. R. Moekti, "Introduction of Superior Grasses as A Part of Regional Partnership Program to Support Livestock Productivity in Bontoharu District within Selayar Island Regency, South Sulawesi, Indonesia", The Business and Management Review, Vol. 10, no. 2, pp. 56-65, 2019.
- [6] A. Rochana, N. P. Indriani, B. Ayuningsih, I. Hernaman, T. Dhalika, D. Rahmat, and S. Suryanah, "Feed Forage and Nutrition Value at Altitudes During the Dry Season in West Java", Animal Production, Vol. 18, no. 2, pp. 85-93, 2016.
- [7] K. Tikam, C. Phatsara, C. Mikled, T. Vearasilp, W. Phunphiphat, J. Chobtang, A. Cherdthong, and K-H Sudekum, "Pangola Grass as Forage for Ruminant Animals: a review", Springerplus, Vol. 2, No. 604, pp. 1-6, 2013. doi: [10.1186/2193-1801-2-604](https://doi.org/10.1186/2193-1801-2-604)
- [8] S. Abd-Aziz, M. Gozan, M. F. Ibrahim, and L-Y. Phang, "Demand and Sustainability of Palm Oil Plantation". In Book: Biorefinery of Oil Producing Plants for Value-Added Products. 2022. DOI: [10.1002/9783527830756.ch2](https://doi.org/10.1002/9783527830756.ch2)
- [9] O. I. Mba, M-J. Dumont, and M. Ngadi, "Palm Oil: Processing, Characterization and Utilization in the Food Industry-A Review. Food Bioscience, Vol. 10, pp. 26-41, 2015. <https://doi.org/10.1016/j.fbio.2015.01.003>
- [10] V. I. Otti, H. I. Ifeanyichukwu, F. C. Nwaorum, and F. U. Ogbuagu, "Sustainable Oil Palm Waste Management in Engineering Development", Civil and Environmental Research, Vol. 6, no. 5, pp. 121-125, 2014.
- [11] T. Astuti, S. A. Akbar, Harissatria, R. M. Sari, D. Afriani, and N. Ropiq, "The Effect of Using Local Microorganism of Ruminant Contents and Crude Enzymes with Different Dosage on The Nutrient Content of Oil Palm Frond. Hasanuddin J. Anim. Sci., Vol. 2, no. 2, pp. 52-57, 2020. DOI: <https://doi.org/10.20956/hajas.v2i2.12212>
- [12] H. Mayulu, "The Nutrient Digestibility of Locally Sheep Fed with Amofer Palm Oil by Product Based Complete Feed, "International Journal of Science and Engineering (IJSE), Vol. 7, no. 2, pp. 106-111, 2014. DOI: <https://doi.org/10.12777/ijse.7.2.106-112>
- [13] H. Maluyu, and S. Suhardi, "The Feed Intake and Daily Weight Gain of Locally Sheep Fed with Amofer Palm Oil Plantation and Mills By Product-Based Complete Feed", International Journal of Science and Engineering (IJSE), Vol. 10, no. 2: 67-73, 2016.
- [14] M. Ishida, and O. H. Hassan, "Utilization of Oil Palm as Cattle Feed", JARQ, Vol. 31, pp. 41-47. 1997.
- [15] A. Rizali, Fahcianto, M. H. Ansari, and A. Wahdi, "Utilization of Waste of Midrib and Palm Oil Leaves Through Fermentation of *Trichoderma sp.* As Beef Cattle Feed", EnviroScientee, Vol. 14, no. 1, pp. 1-7, 2018.

- [16] T. Astuti, "The Effect of Fermented Oil Palm Fronds with Local Microorganism Base on Waste of Livestock on the Content of Fibre Fraction as Ruminant Feeding", Proceeding: Seminar Nasional LPPM University of Jambi, pp. 218-223, 2015.
- [17] R. Sharma, P. Garg, P. Kumar, S. K. Bhatia, and S. Kulshrestha, "Microbial Fermentation and Its Role in Quality Improvement of Fermented Foods", *Fermentation*, Vol. 6, no. 106, pp. 1-20, 2020. doi:10.3390/fermentation6040106
- [18] T. Astuti, Syahro Ali Akbar, Delsi Afrini, M Nasir Rofiq, IrnaHumaira. The Identification of Fungi Colonies Total On The Rumen Content of Cow And Buffalo With Addition of Leaves And Oil Palm Frond. *World Journal of Advanced Research and Reviews*, vol. 8, no. 02, pp 314-317. 2022. DOI: 10.30574/wjarr.2020.8.2.0444
- [19] Statistic Indonesia, "<https://www.bps.go.id/indicator/24/214/1/jumlah-ternak-yang-dipotong-di-rumah-potong-hewan-rph-menurut-provinsi-dan-jenis-ternak.html>", 2021.
- [20] A. G. Williams, and S.E. Withers, "Changes in the Rumen Microbial Population and Its Activities During the Refaunation Period After the Reintroduction of Ciliate Protozoa into the Rumen of Defaunated Sheep", *Can. J. Microbiol.*, Vol. 39, no. 1, pp. 61-69, 1993.
- [21] M.Haq, Shultana Fitra, Sylvia Madusari, and Danielndra Yama. Potensi Kandungan Nutrisi Pakan Berbasis Limbah Pelepah Kelapa Sawit dengan Teknik Fermentasi. Seminar Nasional Sains dan Teknologi 2018. Fakultas Teknik Universitas Muhammadiyah Jakarta , 17 Oktober 2018
- [22] A. Budiansyah, Resmi, Nahrowi, K.G. Wiryawan, M.T. Suhartono, and Y. Widyastuti, "Hidrolisis of Feed Nutrient by Cow Rumen Liquid Enzymes from Slaughterhouse", *AGRINAK*, Vol. 1, No.1, pp. 17-24, 2011.
- [23] W. Pamungkas. Penggunaan Enzim Cairan Rumen Sebagai Alternatif Untuk Mendukung Pemanfaatan Bahan Baku Pakan Ikan Lokal. *Media Akuakultur*. Vol 7, No 1, hal 32-38. 2012. DOI: <http://dx.doi.org/10.15578/ma.7.1.2012.32-38>
- [24] Riswandi, Kualitas Silase Eceng Gondok (*Eichhornia crassipes*) dengan Penambahan Dedak Halus dan Ubi Kayu", *Jurnal Peternakan Sriwijaya*, Vol. 3, no. 1, pp. 1-6, 2014.

# Evaluation of Using Bioactivator

---

## ORIGINALITY REPORT

---

11%	%	11%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** D. Surtina, R.M. Sari, T. Astuti, A.H. Kusuma. "The Effect of Using Some Kind of Manure on the Content of Dry Matter, Organic Matter and Crude Protein of Elephant Grass Cv. Taiwan (*Pennisetum purpureum*) on the First Harvest", IOP Conference Series: Earth and Environmental Science, 2019  
Publication 3%
  - 2** T. Astuti, M. Nasir Rofiq, Nurhaita, U. Santoso. "Analysis of Fibre Fraction of Palm Oil Frond Fermented with Different Microbes and Soluble Carbohydrates Addition as Ruminant Feeding", IOP Conference Series: Earth and Environmental Science, 2019  
Publication 3%
  - 3** AR Gusta, M. Same. "The Effect of Organic Fertilizer and NPK on the Growth of the Master Pepper Plants", IOP Conference Series: Earth and Environmental Science, 2022  
Publication 2%
-



4

Ana Rochana, Nyimas Popi Indriani, Rachmat Wiradimadja, Budi Ayuningsih et al. "Cassava Forage (*Manihot esculenta* Crantz) Ruminant Feed Nutritional Value at Altitudes in the West Java", *Biosciences Biotechnology Research Asia*, 2018

Publication

1 %

5

R Renfiyeni, H Andraini, L Iswaldi. "Growth and yield of *Fragaria* sp. in mixed and volume of plant media", *IOP Conference Series: Earth and Environmental Science*, 2020

Publication

1 %

6

Harissatria, Dara Surtina, Jaswandi, Hendri, T Astuti, D Afrini, R M Sari, J Hendri, A Asri. "The Preservation Epididymis Spermatozoa of Buffalo into Talp Diluent with Addition of Serum Level", *IOP Conference Series: Earth and Environmental Science*, 2021

Publication

1 %

7

Bukola Christianah Adebayo-Tayo, Francis Oluwabunmi Olomitutu, Gboyega Ebenezer Adebami. "Production and evaluation of probioticated mango juice using *Pediococcus pentosaceus* and *Pediococcus acidilactici* during storage at different temperature", *Journal of Agriculture and Food Research*, 2021

Publication

<1 %

8

Diana Nur Afifah, Yesi Pratama Aprilia Ningrum, Tazkiah Syahidah, Nuryanto Nuryanto et al. "Nutrient Content, Organoleptic Quality, and Shelf Life of Sagon Substitute From Lindur (*Bruguiera gymnorrhiza* L.) and Soybean Flour (*Glycine max* L.), as an Alternative Emergency Food", *Frontiers in Nutrition*, 2022

Publication

&lt;1 %

9

Tri Astuti, Delsi Afrini, Alfian Asri, M. Nasir Rofiq. "Identifikasi Morfologi Mikroorganisme Lokal (Mol) Isi Rumen Sapi Dengan Supply Energi Yang Berbeda", *Jurnal Peternakan Mahaputra*, 2021

Publication

&lt;1 %

10

Y Sukaryana, S K Syahpura, Susanti. "Optimization of Addition of Garlic Powder (*Allium sativum*) in Ration to Broiler Performance", *IOP Conference Series: Earth and Environmental Science*, 2022

Publication

&lt;1 %

Exclude quotes  OnExclude matches  < 5 wordsExclude bibliography  On